

Supplementary Table 2. Distribution of serum PFAS in men according to variables: PFOA, PFOS, PFHxS, PFNA, PFDeA

Variable	PFOA			PFOS		
	Low (n=974) ^a	High (n=324)	<i>p</i> -value	Low (n=974)	High (n=324)	<i>p</i> -value
Marital status			0.004 ^b			0.002
Single	166(97.6)	10(2.4)		171(98.0)	5(2.0)	
Married	743(71.8)	292(28.2)		738(75.0)	297(25.0)	
Others	65(74.0)	22(26.0)		65(75.3)	22(24.7)	
Smoking			0.003			0.002
None	263(87.3)	63(12.7)		254(86.4)	72(13.6)	
Ex-smoker	379(70.2)	174(29.8)		382(75.0)	171(25.0)	
Current smoker	332(79.4)	87(20.6)		338(82.2)	81(17.8)	
Hiking boots and sneakers use			0.012			0.027
None	486(81.7)	132(18.3)		465(80.6)	153(19.4)	
< Four times a week	158(69.1)	81(30.9)		171(75.8)	68(24.2)	
≥ Four times a week	330(79.6)	111(20.4)		338(84.3)	103(15.7)	
Coated pot use			0.276			0.294
None	370(79.1)	133(20.9)		363(79.8)	140(20.2)	
< Four times a week	248(80.3)	73(19.7)		251(82.0)	70(18.0)	
≥ Four times a week	356(77.5)	118(22.5)		360(81.8)	114(18.2)	
Coated electric cookers use			0.030			0.008
None	118(73.6)	45(26.4)		121(78.2)	42(21.8)	
< Four times a week	213(82.9)	60(17.1)		219(88.2)	54(11.8)	
≥ Four times a week	643(78.5)	219(21.5)		634(79.5)	228(20.5)	
Coated containers use			0.017			0.011
None	755(77.9)	255(22.1)		750(79.4)	260(20.6)	
< Four times a week	186(79.4)	67(20.6)		193(84.8)	60(15.2)	
≥ Four times a week	33(96.2)	2(3.8)		31(97.1)	4(2.9)	

Coated agent or polish use			0.187			0.277
<Once a week	959(78.7)	320(21.3)		959(81.0)	320(19.0)	
≥Once a week	15(83.8)	4(16.2)		15(86.5)	4(13.5)	
Grilled fish consumption			0.016			0.004
<Once a week	785(79.8)	234(20.2)		788(81.8)	231(18.2)	
≥Once a week	189(75.1)	90(24.9)		186(78.8)	93(21.2)	
Seaweed consumption			0.025			0.001
<Once a week	226(82.8)	61(17.2)		225(86.7)	62(13.3)	
≥Once a week	748(77.6)	263(22.4)		749(79.4)	262(20.6)	

Data are shown as number (%) for categorical variables and as mean ± standard error for continuous variables

BMI: body mass index

a: Unweighted count; b: *p*-value by χ^2 test.