

Supplementary Table 4. Distribution of serum PFAS in women according to variables: PFOA, PFOS, PFHxS, PFNA, PFDeA

Variable	PFOA			PFOS		
	Low (n=1,272) ^a	High (n=423)	<i>p</i> -value	Low (n=1,272)	High (n=423)	<i>p</i> -value
Marital status			0.001 ^b			<0.001
Single	171(98.0)	7(2.0)		169(95.7)	9(4.3)	
Married	931(77.3)	320(22.7)		939(79.1)	312(20.9)	
Others	170(54.8)	96(45.2)		164(58.1)	102(41.9)	
Smoking			0.076			0.015
None	1,195(77.3)	409(22.7)		1,193(78.5)	411(21.5)	
Ex-smoker	38(91.0)	6(9.0)		40(94.5)	4(5.5)	
Current smoker	39(90.5)	8(9.5)		39(92.3)	8(7.7)	
Hiking boots and sneakers use			0.071			0.021
None	792(79.2)	241(20.8)		779(79.7)	254(20.3)	
< Four times a week	226(75.4)	79(24.6)		234(80.9)	71(19.1)	
≥ Four times a week	254(77.0)	103(23.0)		259(76.5)	98(23.5)	
Coated pot use			0.012			0.005
None	497(79.7)	164(20.3)		483(78.0)	178(22.0)	
< Four times a week	332(73.4)	126(26.6)		351(76.8)	107(23.2)	
≥ Four times a week	443(80.0)	133(20.0)		438(82.9)	138(17.1)	
Coated electric cookers use			0.016			0.005
None	149(73.5)	70(26.5)		150(76.9)	69(23.1)	
< Four times a week	315(79.3)	101(20.7)		326(82.4)	90(17.6)	
≥ Four times a week	808(78.6)	252(21.4)		796(78.7)	264(21.3)	
Coated containers use			0.028			0.003
None	964(76.9)	334(23.1)		955(77.9)	343(22.1)	
< Four times a week	273(82.4)	82(17.6)		279(82.2)	76(17.8)	
≥ Four times a week	35(77.9)	7(22.1)		38(96.6)	4(3.4)	

Coated agent or polish use			<0.001			<0.001
<Once a week	1,268(78.2)	422(21.8)		1,269(79.4)	421(20.6)	
≥Once a week	4(38.0)	1(62.0)		3(38.0)	2(62.0)	
Grilled fish consumption			0.129			0.002
<Once a week	1,043(78.5)	315(21.5)		1,042(80.4)	316(19.6)	
≥Once a week	229(76.3)	108(23.7)		230(74.5)	107(25.5)	
Seaweed consumption			<0.001			0.004
<Once a week	334(83.3)	79(16.7)		331(86.1)	82(13.9)	
≥Once a week	938(76.4)	344(23.6)		941(77.1)	341(22.9)	

Data are shown as number (%) for categorical variables and as mean ± standard error for continuous variables

BMI: body mass

a: Unweighted count; b: *p*-value by χ^2 test.